



► Is cannabis safe during preconception, pregnancy and breastfeeding?

CANNABIS EVIDENCE BRIEF



Government
of Canada

Gouvernement
du Canada

Canada 

BRIEFS AVAILABLE IN THIS SERIES:

- ▶ *Is cannabis safe to use? Facts for youth aged 13–17 years.*
- ▶ *Is cannabis safe to use? Facts for young adults aged 18–25 years.*
- ▶ *Does cannabis use increase the risk of developing psychosis or schizophrenia?*
- ▶ *Is cannabis safe during preconception, pregnancy and breastfeeding?*
- ▶ *Is cannabis addictive?*

PURPOSE:

This document provides: key messages for health care providers (HCP) to share with individuals planning a pregnancy, who are pregnant or who are breastfeeding; and information to inform HCPs about the risks and health effects of cannabis use during pregnancy and breastfeeding. Information in this factsheet can be reproduced to develop public education materials, including videos, brochures, pamphlets, posters, etc.

© Her Majesty the Queen in Right of Canada, as represented by the Minister of Health, 2018

Publication date: August 2018

This document may be reproduced in whole or in part for non-commercial purposes, without charge or further permission, provided that it is reproduced for public education purposes and that due diligence is exercised to ensure accuracy of the materials reproduced.

Cat.: H14-264/5-2018E-PDF
ISBN: 978-0-660-27413-3
Pub.: 180234

► **Key messages to share with women planning a pregnancy, and pregnant and breastfeeding women**

Having legal access to cannabis in Canada for people over 18 years old (or 19 in some provinces and territories) does not mean cannabis is safe to use while trying to conceive or while pregnant and/or breastfeeding. As with alcohol or tobacco, cannabis use while trying to conceive or while pregnant and/or breastfeeding carries risks and should be avoided.

General messaging

- Cannabis smoke contains many of the same harmful chemicals as tobacco smoke.¹⁻³
- It is important to tell your healthcare provider if you and/or your partner are using cannabis and you want to become pregnant, you are pregnant or breastfeeding.
- There is no known safe amount of cannabis use during pregnancy or breastfeeding.^{1,4} The safest approach is to not use cannabis.⁵
- If you are using cannabis or cannabis products for medical purposes, talk to your healthcare provider about safer alternatives during pregnancy and breastfeeding.⁵
- If you need more information or support, talk to your healthcare provider.

For individuals planning for pregnancy

- Using cannabis may harm the fertility of both men and women and decrease the ability to become pregnant.^{6,7}

For pregnant women

- Using cannabis during your pregnancy may result in your baby having a lower birth weight.^{1,8}
- Using cannabis during pregnancy may harm your child's brain development and lead to:
 - problems understanding, learning, remembering or succeeding at school
 - hyperactivity, inattentiveness or impulsive behaviour
 - increased risk of depression or anxiety

These effects may last throughout your child's life.^{9,10}

- Using cannabis to help with nausea or vomiting during pregnancy is not recommended. Talk to your healthcare provider about safer alternatives during pregnancy.

Continued on next page

Continued from previous page

For breastfeeding women

- ▶ THC (delta-9-tetrahydrocannabinol), the substance in cannabis responsible for the “high”, is found in the breastmilk of women who smoke cannabis.¹¹ If using cannabis affects your mind and body, it may also affect your child’s mind and body.
- ▶ It is not known whether cannabidiol (CBD), another substance in cannabis, passes into the breastmilk of women using cannabis or CBD-containing products. However like THC, CBD is likely to accumulate in fatty tissues, such as breast tissue.

► Supporting information for health care providers

The Society of Obstetricians and Gynaecologists of Canada recommends that:¹²

- Pregnant women and women of child-bearing age be asked periodically about substance use, including cannabis.
- Healthcare providers should advise pregnant women to abstain from or reduce cannabis use during pregnancy to prevent negative long-term cognitive and behaviour outcomes for exposed children.

Cannabis use among pregnant individuals

More research is needed to clearly understand all the potential effects of cannabis on fertility, pregnant and breastfeeding women, and their offspring.¹

Cannabis is the most frequently used illicit drug during pregnancy.⁹ Although there are no recent Canadian statistics on cannabis use during pregnancy, in a recent (2015) national US survey, approximately 3.5% of pregnant women reported using cannabis during the past month.¹³ Use of cannabis by pregnant women is relatively constant throughout pregnancy: 4.0% of pregnant women use cannabis during their first trimester compared to 3.5% and 2.7% in the second and third trimesters, respectively.¹³

Effects of cannabis on fertility

Using cannabis may decrease the ability to become pregnant:

- In men, using cannabis may decrease the number and the motility of sperm making it harder to get their partner pregnant.⁶
- In women, cannabis may harm fertility through different mechanisms such as menstrual cycle disruptions and decreased egg implantation.^{14, 15} Using cannabis can also decrease the success of an *in vitro* fertilization procedure.⁷

Individuals who are trying to get pregnant should be advised to not use cannabis.

Effect of using cannabis while pregnant

SHORT-TERM EFFECTS ON THE BODY:

- damage blood vessels when smoked¹⁶
- increase heart rate¹
- decrease blood pressure¹⁷

The decrease in blood pressure may cause dizziness or fainting which may cause injury to both mother and fetus as a result of a fall. Women who use cannabis during pregnancy may also be at greater risk of anemia.⁸

SHORT-TERM EFFECTS ON THE BRAIN:

- ▶ sleepiness and fatigue^{18, 19}
- ▶ confusion and forgetfulness¹⁸
- ▶ anxiousness and fear of others (paranoia)¹⁸
- ▶ impaired coordination^{20, 21}

Effects of cannabis on fetuses and breastfed children

During pregnancy, chemicals found in cannabis, such as THC, the substance that produces the “high” and other effects on the brain and body, are carried through the bloodstream to the fetus.²²

Similarly, if cannabis is used while breastfeeding, THC passes from the blood into breastmilk and then on to the baby.¹¹ There may be up to eight times more THC in breast milk than in a mother’s blood.^{23, 24}

THC (and other cannabinoids) are stored in fat²⁵ and are slowly released from the fat back into the bloodstream.²⁶ In heavy users, it can take up to 30 days after stopping cannabis for THC levels to be undetectable in the blood.²⁷ This means that even if one has stopped using cannabis, THC and other cannabinoids may still be in the blood and be passed onto the fetus or the breastfeeding baby.

THC that is passed on from mother to baby during pregnancy and breastfeeding can affect a baby’s health in many ways. For example:

- ▶ Exposure to cannabis during pregnancy is associated with a low birth weight.^{1, 28}
- ▶ Babies exposed to THC through breastmilk may be drowsy, have reduced muscular tone and have poor suckling,²⁹ which could impact breastfeeding success and the baby getting enough breastmilk.

It is possible that the effects may persist depending on the extent and duration of the child’s exposure to cannabis. If cannabis is used during pregnancy or breastfeeding, it may affect a child’s brain development, behaviour and mental health.³⁰ The child may have problems remembering things and paying attention in school and experience difficulty thinking through problems. This could affect their behaviour (e.g. make them hyperactive) and even increase the chance they will use cannabis when they are older.^{9, 10} However, more research is needed to better understand the long-term health effects of cannabis use on fetuses and breastfed babies.

Using cannabis for medical purposes during pregnancy or breastfeeding

The Society of Obstetricians and Gynaecologists of Canada recommends that alternative therapies proven to be safe during preconception, pregnancy and breastfeeding should be recommended over cannabis for medical purposes.³¹

Using cannabis to treat nausea and vomiting of pregnancy

Some women use cannabis to treat nausea and vomiting during their pregnancy.^{32, 33} However, this is not a recommended practice.³⁴ Women are encouraged to ask their healthcare provider about safer alternatives to treat nausea and vomiting during pregnancy.

References

1. National Academies of Sciences, Engineering, and Medicine. The health effects of cannabis and cannabinoids: The current state of evidence and recommendations for research. Washington, D.C.: National Academies Press; 2017.
2. Moir D, Rickert WS, Levasseur G, Larose Y, Maertens R, White P, Desjardins S. A comparison of mainstream and sidestream marijuana and tobacco cigarette smoke produced under two smoking conditions. *Chem Res Toxicol* 2008;21:494-502.
3. Martinasek MP, McGrogan JB, Maysonet A. A systematic review of the respiratory effects of inhalational marijuana. *Respir Care* 2016 Nov;61(11):1543-51.
4. Jaques SC, Kingsbury A, Henshcke P, Chomchai C, Clews S, Falconer J, Abdel-Latif ME, Feller JM, Oei JL. Cannabis, the pregnant woman and her child: Weeding out the myths. *J Perinatol* 2014 Jun;34(6):417-24.
5. Committee opinion summary NO. 722: Marijuana use during pregnancy and lactation. *Obstet Gynecol* 2017 Oct;130(4):931-2.
6. Gundersen TD, Jorgensen N, Andersson AM, Bang AK, Nordkap L, Skakkebaek NE, Priskorn L, Juul A, Jensen TK. Association between use of marijuana and male reproductive hormones and semen quality: A study among 1,215 healthy young men. *Am J Epidemiol* 2015 Sep 15;182(6):473-81.
7. Klonoff-Cohen HS, Natarajan L, Chen RV. A prospective study of the effects of female and male marijuana use on in vitro fertilization (IVF) and gamete intrafallopian transfer (GIFT) outcomes. *Am J Obstet Gynecol* 2006 Feb;194(2):369-76.
8. Gunn JK, Rosales CB, Center KE, Nunez A, Gibson SJ, Christ C, Ehiri JE. Prenatal exposure to cannabis and maternal and child health outcomes: A systematic review and meta-analysis. *BMJ Open* 2016 Apr 5; 6(4):e009986,2015-009986.
9. Porath-Waller AJ. Clearing the smoke on cannabis. maternal cannabis use during pregnancy - an update. Canadian Centre on Substance Abuse; 2015.
10. Grant KS, Petroff R, Isoherranen N, Stella N, Burbacher TM. Cannabis use during pregnancy: Pharmacokinetics and effects on child development. *Pharmacol Ther* 2017 Aug 25.
11. Perez-Reyes M, Wall ME. Presence of delta9-tetrahydrocannabinol in human milk. *N Engl J Med* 1982 09/23;307(0028-4793; 13):819-20.
12. Ordean A, Wong S, Graves L. No. 349-substance use in pregnancy. *J Obstet Gynaecol can* 2017 Oct; 39(10):922,937.e2.
13. Substance Abuse and Mental Health Services Administration. Results from the 2015 national survey on drug use and health: Detailed tables.; 2016.
14. Brents LK. Marijuana, the endocannabinoid system and the female reproductive system. *Yale J Biol Med* 2016 Jun 27;89(2):175-91.
15. Sharma R, Biedenharn KR, Fedor JM, Agarwal A. Lifestyle factors and reproductive health: Taking control of your fertility. *Reprod Biol Endocrinol* 2013 Jul 16;11:66,7827-11-66.

16. Wang X, Derakhshandeh R, Liu J, Narayan S, Nabavizadeh P, Le S, Danforth OM, Pinnamaneni K, Rodriguez HJ, Luu E, *et al.* One minute of marijuana secondhand smoke exposure substantially impairs vascular endothelial function. *J Am Heart Assoc* 2016 Jul 27;5(8):10.1161/JAHA.116.003858.
17. Benowitz NL, Jones RT. Cardiovascular and metabolic considerations in prolonged cannabinoid administration in man. *J Clin Pharmacol* 1981 Aug-Sep;21(8-9 Suppl):214S-23S.
18. World Health Organization (WHO). The health and social effects of nonmedical cannabis use.
19. Whiting PF, Wolff RF, Deshpande S, Di Nisio M, Duffy S, Hernandez AV, Keurentjes JC, Lang S, Misso K, Ryder S, *et al.* Cannabinoids for medical use: A systematic review and meta-analysis. *JAMA* 2015 Jun 23-30;313(24):2456-73.
20. Hartman RL, Huestis MA. Cannabis effects on driving skills. *Clin Chem* 2013 Mar;59(3):478-92.
21. Weinstein A, Brickner O, Lerman H, Greemland M, Bloch M, Lester H, Chisin R, Sarne Y, Mechoulam R, Bar-Hamburger R, *et al.* A study investigating the acute dose-response effects of 13 mg and 17 mg delta 9- tetrahydrocannabinol on cognitive-motor skills, subjective and autonomic measures in regular users of marijuana. *J Psychopharmacol* 2008 Jun;22(4):441-51.
22. Bar-Oz B, Klein J, Karaskov T, Koren G. Comparison of meconium and neonatal hair analysis for detection of gestational exposure to drugs of abuse. *Arch Dis Child Fetal Neonatal Ed* 2003 Mar;88(2):F98-F100.
23. Alpar A, Di Marzo V, Harkany T. At the tip of an iceberg: Prenatal marijuana and its possible relation to neuropsychiatric outcome in the offspring. *Biol Psychiatry* 2016 Apr 1;79(7):e33-45.
24. Perez-Reyes M. Marijuana smoking: Factors that influence the bioavailability of tetrahydrocannabinol. *NIDA Res Monogr* 1990;99(1046-9516):42-62.
25. Huestis MA. Human cannabinoid pharmacokinetics. *Chem.Biodivers.* 2007 08;4(1612-1880; 1612-1872; 8):1770-804.
26. Karschner EL, Swortwood MJ, Hirvonen J, Goodwin RS, Bosker WM, Ramaekers JG, Huestis MA. Extended plasma cannabinoid excretion in chronic frequent cannabis smokers during sustained abstinence and correlation with psychomotor performance. *Drug Test Anal* 2015 Jun 11.
27. Bergamaschi MM, Karschner EL, Goodwin RS, Scheidweiler KB, Hirvonen J, Queiroz RH, Huestis MA. Impact of prolonged cannabinoid excretion in chronic daily cannabis smokers' blood on per se drugged driving laws. *Clin Chem* 2013 Mar;59(3):519-26.
28. Grant P, Gandhi P. A case of cannabis-induced pancreatitis. *JOP.* 2004 01;5(1590-8577; 1590-8577; 1):41-3.
29. Liston J. Breastfeeding and the use of recreational drugs--alcohol, caffeine, nicotine and marijuana. *Breastfeed Rev* 1998 Aug;6(2):27-30.
30. Astley SJ, Little RE. Maternal marijuana use during lactation and infant development at one year. *Neurotoxicol Teratol* 1990 03;12(0892-0362; 2):161-8.
31. SOGC Position Statement: Marijuana Use during Pregnancy [Internet]; c2017. Available from: <https://sogc.org/files/letSOGCstatementCannabisUse.pdf>.
32. Westfall RE, Janssen PA, Lucas P, Capler R. Survey of medicinal cannabis use among childbearing women: Patterns of its use in pregnancy and retroactive self-assessment of its efficacy against 'morning sickness'. *Complement Ther Clin Pract* 2006 Feb;12(1):27-33.
33. Roberson EK, Patrick WK, Hurwitz EL. Marijuana use and maternal experiences of severe nausea during pregnancy in hawaii. *Hawaii J Med Public Health* 2014 Sep;73(9):283-7.
34. Volkow ND, Compton WM, Wargo EM. The risks of marijuana use during pregnancy. *Journal of American Medical Association* 2016:E1-2.